









































- Melzer, I., Melzer, I., Kurz, I., Shahar, D., & Oddsson, L. I. E. (2010). Do voluntary step reactions in dual task conditions have an added value over single task for fall prediction? A prospective study. *Aging Clin Exp Res*, 22(5), 6–360.
- Grahn, J. A., & Brett, M. (2007). Rhythm and beat perception in motor areas of the brain. *Journal of cognitive neuroscience*, 19(5), 893-906.
- Vasudevan, E. V., & Bastian, A. J. (2009). Split-belt treadmill adaptation shows different functional networks for fast and slow human walking. *Journal of Neurophysiology*, 102(1), 183-191.
- Choi, J. T., & Bastian, A. J. (2007). Adaptation reveals independent control networks for human walking. *Nature neuroscience*, 10(8), 1055-1062.
- Reisman, D. S., Wityk, R., Singer, K., & Bastian, A. J. (2009). Split-belt treadmill adaptation transfers to overground walking in persons poststroke. *Neurorehabilitation and neural repair*, 23(7), 735-744.
- Thompson, T. W., Waskom, M. L., & Gabrieli, J. D. (2016). Intensive working memory training produces functional changes in large-scale frontoparietal networks. *Journal of cognitive neuroscience*.
- Bherer, L., Kramer, A. F., Peterson, M. S., Colcombe, S., Erickson, K., & Bécic, E. (2005). Training effects on dual-task performance: are there age-related differences in plasticity of attentional control?. *Psychology and aging*, 20(4), 695.
- Shigematsu, R., & Okura, T. (2006). A novel exercise for improving lower-extremity functional fitness in the elderly. *Aging clinical and experimental research*, 18(3), 242-248.